

$$\langle x, y \rangle \leq \|x\| \|y\|$$

$$\frac{d\vec{v}}{dt} = \vec{a} \quad \frac{d\vec{x}}{dt} = \vec{v}$$

$$d\vec{v} = \vec{a} dt \quad d\vec{x} = (\vec{v}_0 + \vec{a}t) dt$$

$$\int d\vec{v} = \int \vec{a} dt \quad \int d\vec{x} = (\vec{v}_0 + \vec{a}t) dt$$

$$\vec{v} = \vec{v}_0 + \vec{a}t \quad \int d\vec{x} = (\vec{v}_0 + \vec{a}t) dt$$

$$\vec{x} = \vec{x}_0 + \vec{v}_0 t + \frac{1}{2} \vec{a} t^2$$



$$\hat{H}|\psi_n(t)\rangle = i\hbar \frac{\partial}{\partial t} |\psi_n(t)\rangle$$

$$\frac{1}{c^2} \frac{\partial^2 \phi_n}{\partial t^2} - \nabla^2 \phi_n + \left(\frac{mc}{\hbar}\right)^2 \phi_n = 0$$

$$\hbar \frac{\partial}{\partial t_0} s = s / \hbar \frac{\partial}{\partial t_1} s = p_i o s, i=1, \dots, k.$$

$$f(Q_i) = \sum_{d_i=1}^{\infty} \frac{(2d_i-1)!}{(d_i!)^2} Q_i^{d_i}$$

$$d(x, z) \leq d(x, y) + d(y, z)$$

$$\frac{d\vec{v}}{dt} = \vec{a} \quad \frac{d\vec{x}}{dt} = \vec{v}$$

$$d\vec{v} = \vec{a} dt \quad d\vec{x} = (\vec{v}_0 + \vec{a}t) dt$$

$$\int d\vec{v} = \int \vec{a} dt \quad \int d\vec{x} = (\vec{v}_0 + \vec{a}t) dt$$

$$\vec{v} = \vec{v}_0 + \vec{a}t \quad \int d\vec{x} = (\vec{v}_0 + \vec{a}t) dt$$

$$\vec{x} = \vec{x}_0 + \vec{v}_0 t + \frac{1}{2} \vec{a} t^2$$

$$\hat{H}|\psi_n(t)\rangle = i\hbar \frac{\partial}{\partial t} |\psi_n(t)\rangle$$

$$\frac{1}{c^2} \frac{\partial^2 \phi_n}{\partial t^2} - \nabla^2 \phi_n + \left(\frac{mc}{\hbar}\right)^2 \phi_n = 0$$

$$\hbar \frac{\partial}{\partial t_0} s = s / \hbar \frac{\partial}{\partial t_1} s = p_i o s, i=1, \dots, k.$$

$$f(Q_i) = \sum_{d_i=1}^{\infty} \frac{(2d_i-1)!}{(d_i!)^2} Q_i^{d_i}$$

$$d(x, z) \leq d(x, y) + d(y, z)$$

$$\langle x, y \rangle \leq \|x\| \|y\|$$

$$\frac{d\vec{v}}{dt} = \vec{a} \quad \frac{d\vec{x}}{dt} = \vec{v}$$

$$d\vec{v} = \vec{a} dt \quad d\vec{x} = (\vec{v}_0 + \vec{a}t) dt$$

$$\int d\vec{v} = \int \vec{a} dt \quad \int d\vec{x} = (\vec{v}_0 + \vec{a}t) dt$$

$$\vec{v} = \vec{v}_0 + \vec{a}t \quad \int d\vec{x} = (\vec{v}_0 + \vec{a}t) dt$$

$$\vec{x} = \vec{x}_0 + \vec{v}_0 t + \frac{1}{2} \vec{a} t^2$$

$$\langle x, y \rangle \leq \|x\| \|y\|$$

$$\frac{d\vec{v}}{dt} = \vec{a} \quad \frac{d\vec{x}}{dt} = \vec{v}$$

$$d\vec{v} = \vec{a} dt \quad d\vec{x} = (\vec{v}_0 + \vec{a}t) dt$$

$$\int d\vec{v} = \int \vec{a} dt \quad \int d\vec{x} = (\vec{v}_0 + \vec{a}t) dt$$

$$\vec{v} = \vec{v}_0 + \vec{a}t \quad \int d\vec{x} = (\vec{v}_0 + \vec{a}t) dt$$

$$\vec{x} = \vec{x}_0 + \vec{v}_0 t + \frac{1}{2} \vec{a} t^2$$

$$\hat{H}|\psi_n(t)\rangle = i\hbar \frac{\partial}{\partial t} |\psi_n(t)\rangle$$

$$\frac{1}{c^2} \frac{\partial^2 \phi_n}{\partial t^2} - \nabla^2 \phi_n + \left(\frac{mc}{\hbar}\right)^2 \phi_n = 0$$

$$\hbar \frac{\partial}{\partial t_0} s = s / \hbar \frac{\partial}{\partial t_1} s = p_i o s, i=1, \dots, k.$$

$$f(Q_i) = \sum_{d_i=1}^{\infty} \frac{(2d_i-1)!}{(d_i!)^2} Q_i^{d_i}$$

$$d(x, z) \leq d(x, y) + d(y, z)$$